

Exercise Prescription: Resistance Training for Patients with DM and NWB on Lower Extremity

 APTA Academy of Clinical Electrophysiology & Wound Management.	Prescription Component	Resistance Training	
F	Frequency 	<ul style="list-style-type: none"> 2-3 days/week with at least 48 hours between training the same muscle group^{1,2} 	
I	Intensity 	<ul style="list-style-type: none"> Moderate (50-69% of 1-RM)¹ or Vigorous (70-85% of 1-RM)¹ 	
T	Time 	<ul style="list-style-type: none"> 10 – 15 repetitions per set^{1,2} 1 – 3 sets per exercise type^{1,2} 	
T	Type 	<ul style="list-style-type: none"> Seated resistance using:^{1,2} <ul style="list-style-type: none"> Body weight Elastic bands Free weights Weight machines Work major muscle groups with 8-10 exercises¹ 	
V	Volume 	<ul style="list-style-type: none"> Total of repetitions and sets 	
P	Progression 	<ul style="list-style-type: none"> Can increase resistance (intensity) Increase number of repetitions or sets Increase frequency of resistance training 	
Why?	Reasoning 	<ul style="list-style-type: none"> With strength training, muscle needs rest prior to repeated bouts, just like for people without diabetes. Strength training supports improved diabetes management. 	
Benefits¹⁻³		<ul style="list-style-type: none"> Better glycemic control Improved insulin sensitivity Lower blood pressure Lower A1C Improved lipid profile Improved function Decreased depression Improved quality of life 	

Modified from Table 2 Kanaley JA, et al. 2022² and consistent with Standards of Medicine in Diabetes.³ NWB: Non-weight-bearing; RM: repetition maximum;

Before performing resistance training:

1. Check blood glucose
2. Check blood pressure
3. Consider orthopedic limitations

Following exercise:

1. Check blood glucose
2. Check blood pressure
3. Remind patient to wait 48 hours to lift again

Example Exercise Prescription Related to Non-weight bearing Resistance Exercise:

Resistance training for each of the major muscle groups (e.g., seated arm curls with 10-lb weight when participant has a 20-lb 1-RM for arm curl) (**type** and **intensity**) for 12 repetitions per set x 2 sets (**time**) x 2 times per week (**frequency**). The **volume** of this exercise would be 4 sets of 12 repetitions per week. This activity could be progressed by increasing number of repetitions (e.g., 15 repetitions per set) OR number of sets (e.g., change from 2 to 3) OR frequency of bouts (e.g., 3 times per week) OR intensity of bouts (e.g., moving from moderate (50-69% of 1-RM) to vigorous (70-85% of 1-RM)) (**progression**).

References:

1. ACSM. ACSM's guidelines for exercise testing and prescription. 12th ed. Philadelphia, Wolters Kluwer, 2026.
2. Kanaley JA, Colberg SR, Corcoran MH, et al. Exercise/physical activity in individuals with type 2 diabetes: a consensus statement from the American College of Sports Medicine. *Med Sci Sports Exerc.* 2022;54(2):353-368.
3. American Diabetes Association Professional Practice Committee. 5. Facilitating positive health behaviors and well-being to improve health outcomes: standards of care in diabetes—2025. *Diabetes Care.* 2025;48(Suppl. 1):S86-S127.

Before performing resistance training:

1. Check blood glucose
2. Check blood pressure
3. Consider orthopedic limitations

Following exercise:

1. Check blood glucose
2. Check blood pressure
3. Remind patient to wait 48 hours to lift again